

Oregon Statewide Snapshot

February 24, 2020

Comagine Health continually tracks Oregon performance on a number of quality and resource use metrics. The table on the following pages provides a snapshot of Oregon's overall primary care clinic performance. The table shows Oregon's score for each measure compared to local and national benchmarks. It also includes information on the three year trend for each measure.

For the local benchmark, Comagine Health calculates the Oregon Achievable Benchmark of Care (ABC), labeled Oregon's Best Benchmark. This benchmark provides a method to identify performance levels already being achieved by "best-in-class" primary care clinics within Oregon. "Best-in-class" Oregon clinic scores are calculated using the paired mean rate of the highest performing clinics providing care to at least 10 percent of the patient population.

The national benchmarks come from National Committee for Quality Assurance (NCQA) HEDIS® (Healthcare Effectiveness Data and Information Set). For each measure, the national mean and 90th percentile were calculated by weighting the 2019 HEDIS® benchmarks for each line of business based on the proportion of each population (commercial, Medicaid and Medicare).

For more information on the measure calculations, please refer to the Technical Appendix that can be found here:

<http://q-corp.org/reports/technical-appendix>

Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average
				■ 2016 ■ 2017 ■ 2018
Women's Screenings				
Breast Cancer Screening ‡	62.0%	85.0%	79.7%	64.6% (2016) 63.7% (2017) 62.0% (2018)
Cervical Cancer Screening	60.5%	83.0%	75.9%	57.4% (2016) 57.6% (2017) 60.5% (2018)
Chlamydia Screening	48.8%	70.0%	65.8%	42.2% (2016) 45.0% (2017) 48.8% (2018)
				■ 2016 ■ 2017 ■ 2018
Adult Quality Measures				
Antidepressant Medication Management (Short Term)	56.3%	77.0%	68.3%	55.2% (2016) 53.9% (2017) 56.3% (2018)
Antidepressant Medication Management (Long Term)	43.4%	65.0%	53.1%	36.0% (2016) 41.3% (2017) 43.4% (2018)
Admissions for Ambulatory Sensitive Conditions - Overall per 100,000 † ‡	1,032.7	92.9	N/A	724.1 (2016) 705.3 (2017) 626.7 (2018)
Admissions for Ambulatory Sensitive Conditions - Acute per 100,000 † ‡	308.6	9.4	N/A	308.6 (2016) 318.8 (2017) 382.4 (2018)
Admissions for Ambulatory Sensitive Conditions - Chronic per 100,000 † ‡	724.1	54.2	N/A	1,032.7 (2016) 1,024.1 (2017) 1,009.0 (2018)

Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average
Pediatric Quality Measures				■ 2016 ■ 2017 ■ 2018
Well-Child Visits in the First 15 Months of Life, Six or More	68.1%	86.0%	75.9%	
Well-Child Visits in the 3rd, 4th, 5th and 6th Years of Life	65.9%	86.0%	82.3%	
Adolescent Well-Care Visits (age 12-21)	40.3%	64.0%	63.7%	
Developmental Screenings in 1st, 2nd and 3rd Year of Life	66.6%	89.0%	N/A	
Diabetes Management				■ 2016 ■ 2017 ■ 2018
Diabetes Blood Sugar (HbA1c) Screening ‡ **	84.8%	95.0%	95.1%	
Diabetes Kidney Disease Monitoring ‡ **	72.1%	94.0%	95.4%	

Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average		
				2016	2017	2018
Use of Services				2016	2017	2018
30-Day All-Cause Readmissions, Unadjusted † ‡	10.8%	5.0%	N/A	9.3%	9.8%	10.8%
Potentially Avoidable Emergency Department Visits, Percentage † ‡	11.1%	7.0%	N/A	11.3%	10.8%	11.1%
Potentially Avoidable Emergency Department Visits, Rate per 1,000 † ‡	36.2	10.9	N/A	40.9	36.7	36.2
Generic Prescription Fills, Antihypertensives	99.5%	100.0%	N/A	98.6%	99.2%	99.5%
Generic Prescriptions Fills, SSRIs	99.2%	100.0%	N/A	98.6%	99.0%	99.2%
Generic Prescriptions Fills, Statins	99.4%	100.0%	N/A	96.7%	99.2%	99.4%
Emergency Department Visits per 1,000 † ‡	311.2	143.0	305.5	350.7	328.6	311.2
Outpatient Visits per 1,000 ‡	3,979.5	N/A	N/A	4173.4	4068.8	3979.5

Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average
				■ 2016 ■ 2017 ■ 2018
Appropriate Care				
Appropriate Low Back Pain Imaging (age 18-50)	82.3%	93.0%	80.7%	81.9% 82.7% 82.3%
Appropriate Testing for Children with Pharyngitis (age 2-18)	86.8%	99.0%	89.6%	79.9% 84.1% 86.8%
Appropriate Treatment for Children With Upper Respiratory Infection	98.9%	100.0%	95.2%	96.6% 98.4% 98.9%
Avoidance of Antibiotic Treatment in Adults With Acute Bronchitis	52.7%	77.0%	45.2%	38.6% 44.6% 52.7%

* Benchmarks use a weighted formula based on the proportion of commercial, Medicaid and Medicare members for each measure

** This measure was not reported in 2017

† Lower scores indicate higher quality

‡ Measure includes Medicare Fee-For-Service data