

Oregon Statewide Snapshot

April 13, 2021

Comagine Health continually tracks Oregon performance on a number of quality and resource use metrics. The table on the following pages provides a snapshot of Oregon's overall primary care clinic performance. The table shows Oregon's score for each measure compared to local and national benchmarks. It also includes information on the three year trend for each measure.

For the local benchmark, Comagine calculates the Oregon Achievable Benchmark of Care (ABC), labeled Oregon's Best Benchmark. This benchmark provides a method to identify performance levels already being achieved by "best-in-class" primary care clinics within Oregon. "Best-in-class" Oregon clinic scores are calculated using the paired mean rate of the highest performing clinics providing care to at least 10 percent of the patient population.

The national benchmarks come from National Committee for Quality Assurance (NCQA) HEDIS® (Healthcare Effectiveness Data and Information Set). For each measure, the national mean and 90th percentile were calculated by weighting the 2020 Report Year HEDIS® benchmarks for each line of business based on the proportion of each population (commercial, Medicaid and Medicare).

For more information on the measure calculations, please refer to the Technical Appendix that can be found here:

http://q-corp.org/reports/technical-appendix



Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average
Women's Screenings				■ 2017 ■ 2018 ■ 2019
Breast Cancer Screening ‡	62.7%	85.3%	75.8%	63.7% 62.0% 62.7%
Cervical Cancer Screening	60.2%	81.2%	76.9%	57.6% 60.5% 60.2%
Chlamydia Screening	48.7%	67.7%	66.3%	45.0% 48.8% 48.7%
Adult Quality Measures				■ 2017 ■ 2018 ■ 2019
Antidepressant Medication Management (Short Term)	63.6%	78.6%	68.7%	53.9% 56.3% 63.6%
Antidepressant Medication Management (Long Term)	48.6%	64.6%	53.0%	41.3% 43.4% 48.6%
Admissions for Ambulatory Sensitive Conditions - Chronic per 100,000 † ‡	724.1	233.1	N/A	724.1 724.1 705.3
Admissions for Ambulatory Sensitive Conditions - Acute per 100,000 † ‡	297.9	43.6	N/A	297.9 308.6 318.8
Admissions for Ambulatory Sensitive Conditions - Overall per 100,000 † ‡	1,022.0	123.6	N/A	1,022.0 1,032.7 1,024.1



Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average
Pediatric Quality Measures				■ 2017 ■ 2018 ■ 2019
Well-Child Visits in the First 15 Months of Life, Six or More	72.6%	89.5%	78.9%	61.8% 68.1% 72.6%
Well-Child Visits in the 3rd, 4th, 5th and 6th Years of Life	71.4%	90.1%	83.7%	65.6% 65.9% 71.4%
Adolescent Well-Care Visits (age 12-21)	44.0%	70.6%	65.4%	40.1% 40.3% 44.0%
Developmental Screenings in 1st, 2nd and 3rd Year of Life	70.3%	91.3%	N/A	62.3% 66.6% 70.3%
Diabetes Management				■ 2018 ■ 2019
Diabetes Blood Sugar (HbA1c) Screening ‡ **	87.8%	94.9%	93.1%	85.3% 87.8%
Diabetes Kidney Disease Monitoring ‡ **	82.6%	93.1%	92.5%	81.6% 82.6%



Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Tre	nd for Ore	egon Average
Use of Services				■ 2017	■ 2018	■ 2019
30-Day All-Cause Readmissions, Unadjusted†‡	11.4%	5.8%	N/A	9.8% 10.8% 11.4%		
Potentially Avoidable Emergency Department Visits, Percentage † ‡	10.8%	7.0%	N/A	10.8% 11.1% 10.8%		
Potentially Avoidable Emergency Department Visits, Rate per 1,000 † ‡	36.7	14.7	N/A		36.7 36.2 36.7	
Generic Prescription Fills, Antihypertensives	99.7%	100.0%	N/A			99.2% 99.5% 99.7%
Generic Prescriptions Fills, SSRIs	99.3%	100.0%	N/A			99.0% 99.2% 99.3%
Generic Prescriptions Fills, Statins	99.3%	100.0%	N/A			99.2% 99.4% 99.3%
Emergency Department Visits per 1,000 † ‡	315.4	171.7	N/A	328.6 311.2 315.4		
Outpatient Visits per 1,000 ‡	4,173.8	N/A	N/A			4068.8 3979.5 4173.8



Measure	Oregon Average	Oregon's Best Benchmark	Combined HEDIS National 90th Percentile*	3-Year Trend for Oregon Average
Appropriate Care				■ 2017 ■ 2018 ■ 2019
Appropriate Low Back Pain Imaging (age 18-50)	80.9%	91.1%	82.4%	82.7% 82.3% 80.9%
Appropriate Testing for Pharyngitis (age 3 years and older) ***	83.3%	96.1%	85.1%	83.3%
Appropriate Treatment for Upper Respiratory Infection (age 3 months and older) ***	92.3%	98.3%	90.2%	92.3%
Avoidance of Antibiotic Treatment for Acute Bronchiolitis/Bronchitis (age 3 months and older) ***	50.8%	80.3%	56.4%	50.8%

^{*} Benchmarks use a weighted formula based on the proportion of commercial, Medicaid and Medicare members for each measure

^{**} This measure was not reported in 2017

^{***} The age range for this measure was expanded in 2019; 2017 and 2018 data is not reported

[†] Lower scores indicate higher quality

[‡] Measure includes Medicare Fee-For-Service data